


A MATHEMATICAL TRIBUTE
to Professor
José María Montesinos Amilibia



DEPARTAMENTO DE GEOMETRÍA Y TOPOLOGÍA
FACULTAD DE CIENCIAS MATEMÁTICAS – UCM



**A MATHEMATICAL TRIBUTE
to Professor José María Montesinos Amilibia**

© Los autores de las comunicaciones

Edita

Departamento de Geometría y Topología
Facultad de Ciencias Matemáticas – UCM
Plaza de Ciencias, 3
Ciudad Universitaria
28040 Madrid

Cubierta

Raquel Díaz
Tetraedro inscrito en un cubo
(Dóblese por las líneas negras finas, córtese por las líneas negras gruesas.)

ISBN: 978-84-608-1684-3
Depósito Legal: M-2056-2016

Imprenta

Ulzama Digital

Impreso en España



Professor José María Montesinos Amilibia

Oh my knots!

This volume contains the contributions presented by several colleagues as a tribute to the mathematical and human qualities of José María Montesinos Amilibia on the occasion of his seventieth birthday. The editors would like to express their thanks to the contributors and their very especial gratitude to José María for his example through many years of scientific and personal contact.

*Marco Castrillón
Elena Martín-Peinador
José M. Rodríguez-Sanjurjo
Jesús M. Ruiz*

Contents

Lectures addressed on the 8 of September, 2015

Morphismes analytiques finis et revêtements ramifiés.....	1
<i>Claude Weber</i>	
Algunas contribuciones matemáticas del Profesor José María Montesinos Amilibia.....	23
<i>Maite Lozano</i>	
José María Montesinos Amilibia. Biographical Sketches.....	33
<i>Sebastià Xambó</i>	

Contributions on the occasion of JMMA's 70 anniversary

Proof of the rigidity of model nilpotent Lie algebras by means of the Internal Set Theory.....	63
<i>J. M. Ancochea Bermúdez, R. Campoamor Stursberg</i>	
Una caracterización de las proyecciones de Lagrange.....	75
<i>Jorge L. Andrés, Jesús Otero</i>	
The dodecahedron: from intersections of quadrics to Borromean rings.....	85
<i>Enrique Artal Bartolo, Santiago López de Medrano, María Teresa Lozano</i>	
ρ -pairs in graphs representing surfaces.....	105
<i>P. Bandieri</i>	
Fixed points, bounded orbits and attractors of planar flows.....	125
<i>Héctor Barge, José M. R. Sanjurjo</i>	
The contact structure in the space of light rays.....	133
<i>Alfredo Bautista, Alberto Ibort, Javier Lafuente</i>	
Universal groups and super regular tessellations.....	161
<i>G. Brumfiel, H. Hilden, M. T. Lozano, J. M. Montesinos-Amilibia, E. Ramirez, H. Short, D. Tejada, M. Toro</i>	
Groups of automorphisms of bordered 2-tori.....	181
<i>Emilio Bujalance, José Javier Etayo, Ernesto Martínez</i>	

Classifying PL 4-manifolds via crystallizations: results and open problems ...	199
<i>Maria Rita Casali, Paola Cristofori, Carlo Gagliardi</i>	
On certain classes of closed 3-manifolds with different geometric structures...	227
<i>Alberto Cavicchioli, Fulvia Spaggiari</i>	
Algunas exploraciones matemáticas del mundo.....	243
<i>Capi Corrales Rodrigáñez</i>	
A note on regular branched foldings	259
<i>Antonio F. Costa</i>	
Partially flat surfaces solving k -Hessian perturbed equations	265
<i>Gregorio Díaz, Jesús Ildelfonso Díaz</i>	
Hyperbolic surfaces of genus 3 with symmetry \mathfrak{S}_4	295
<i>Raquel Díaz, Victor González Aguilera</i>	
Modelo plano conforme del Plano Proyectivo	307
<i>Fernando Etayo, Ujué R. Trías</i>	
Toroidal Dehn Surgeries	317
<i>Mario Eudave-Muñoz, Enrique Ramírez-Losada</i>	
The open quadrant problem: A topological proof	337
<i>José F. Fernando, J. M. Gamboa, Carlos Ueno</i>	
A la búsqueda de la espiritualidad perdida. Meditar itinerante acerca del número, el tacto, la duración y el Arte so pretexto de las Matemáticas y la matematización del mundo	351
<i>Jesús Fortea Pérez</i>	
Persistencia uniforme de atractores	373
<i>Antonio Giraldo, Víctor Fernández Laguna, José M. Rodríguez Sanjurjo</i>	
2-dimensional stratifolds.....	395
<i>J. C. Gómez-Larrañaga, F. González-Acuña, Wolfgang Heil</i>	
Surface knot groups and 3-manifold groups	407
<i>Francisco González-Acuña, Arturo Ramírez</i>	
Some geometric properties of variable exponent Lebesgue spaces	415
<i>Francisco L. Hernández, Cesar Ruiz</i>	
Carousel wild knots are ambient homogeneous	423
<i>Gabriela Hinojosa, Cynthia Verjovsky Marcotte, Alberto Verjovsky</i>	
Global bifurcation for Fredholm operators	437
<i>Julián López-Gómez</i>	

Pascal's triangle, Stirling numbers and the Euler characteristic.....	453
<i>Ana Luzón, Manuel A. Morón, Felipe Prieto-Martínez</i>	
Probabilidades en espacios topológicos.....	463
<i>Juan Margalef-Roig, Salvador Miret-Artés, Enrique Outerelo-Domínguez</i>	
Compact Hausdorff group topologies for the additive group of real numbers ..	491
<i>Elena Martín-Peinador, Monserrat Bruguera Padró</i>	
Unraveling the Dogbone Space.....	499
<i>Diego Mondéjar Ruiz</i>	
$SL(3, \mathbb{C})$ -character variety of torus knots	515
<i>Vicente Muñoz, Jonathan Sánchez</i>	
Spectral limits of semiclassical commuting self-adjoint operators	527
<i>Álvaro Pelayo, S. Vũ Ngọc</i>	
Nontrivial twisted Alexander polynomials.....	547
<i>Joan Porti</i>	
Matrices de rotaciones, simetrías y roto-simetrías	559
<i>María Jesús de la Puente</i>	
La Biblioteca de El Escorial. Un culto a la matemática	565
<i>María Concepción Romo Santos</i>	
Geometría en el siglo XIV: los trabajos de Thomas Bradwardine.....	571
<i>Juan Tarrés Freixenet</i>	

The open quadrant problem: A topological proof

José F. FERNANDO*, J.M. GAMBOA† and Carlos UENO‡

Departamento de Álgebra
Facultad de Ciencias Matemáticas
Universidad Complutense de Madrid
28040 Madrid, Spain
josefer@mat.ucm.es jmgamboa@mat.ucm.es

Dipartimento di Matematica
Università degli studi di Pisa
56127 Pisa, Italy
cuenjac@mail.dm.unipi.it

Dedicated to José María Montesinos on the occasion of his 70th birthday.

ABSTRACT

In this work we present a new polynomial map $f := (f_1, f_2) : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ whose image is the open quadrant $\mathcal{Q} := \{x > 0, y > 0\} \subset \mathbb{R}^2$. The proof of this fact involves arguments of topological nature that avoid hard computer calculations. In addition each polynomial $f_i \in \mathbb{R}[x, y]$ has degree ≤ 16 and only 11 monomials, becoming the simplest known map solving the open quadrant problem.

2010 Mathematics Subject Classification: 14P10, 26C99, 52A10.

Key words: Polynomial map, polynomial image, semialgebraic set, open quadrant, total degree, total number of monomials.

*Author supported by Spanish GAAR MTM2011-22435, Grupos UCM 910444 and the “National Group for Algebraic and Geometric Structures, and their Applications” (GNASA - INdAM). His one year research stay in the Dipartimento di Matematica of the Università di Pisa is partially supported by MECD grant PRX14/00016.

†Author supported by Spanish GAAR MTM2011-22435 and Grupos UCM 910444.

‡Author supported by ‘Scuola Galileo Galilei’ Research Grant at the Dipartimento di Matematica of the Università di Pisa and Spanish GAAR MTM2011-22435.

1. Introduction

Although it is usually said that the first work in Real Geometry is due to Harnack [13], who obtained an upper bound for the number of connected components of a non-singular real algebraic curve in terms of its genus, modern Real Algebraic Geometry was born with Tarski's article [15], where it is proved that the image of a semialgebraic set under a polynomial map is a semialgebraic set. We are interested in studying what might be called the 'inverse problem'. In the 1990 *Oberwolfach Reelle algebraische Geometrie* week [12] the second author proposed:

Problem 1.1 *Characterize the (semialgebraic) subsets of \mathbb{R}^m that are either polynomial or regular images of \mathbb{R}^n .*

A map $f := (f_1, \dots, f_m) : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a *polynomial map* if its components $f_k \in \mathbb{R}[\mathbf{x}] := \mathbb{R}[x_1, \dots, x_n]$ are polynomials. Analogously, f is a *regular map* if its components can be represented as quotients $f_k = \frac{g_k}{h_k}$ of two polynomials $g_k, h_k \in \mathbb{R}[\mathbf{x}]$ such that h_k never vanishes on \mathbb{R}^n . A subset $S \subset \mathbb{R}^n$ is *semialgebraic* when it admits a description by a finite boolean combination of polynomial equalities and inequalities.

Open semialgebraic sets deserve a special attention in connection with the real Jacobian Conjecture [14]. In particular the second author stated in [12] the 'open quadrant problem':

Problem 1.2 *Determine whether the open quadrant $\Omega := \{x > 0, y > 0\}$ of \mathbb{R}^2 is a polynomial image of \mathbb{R}^2 .*

This problem stimulated the interest of many specialists in the field. However, only after twelve years a first solution was found in [4] and presented by the first author in the 2002 *Oberwolfach Reelle algebraische Geometrie* week [2].

The open quadrant problem was the germ of a more systematic study of 'Polynomial and regular images of Euclidean spaces' developed by the authors during the last decade and which was the topic of the Ph.D. Thesis of the third author [16]. Since then we have worked on this issue with two main objectives:

- Finding obstructions to be an either polynomial or regular image.
- Proving (constructively) that large families of semialgebraic sets with piecewise linear boundary (convex polyhedra, their interiors, complements and the interiors of their complements) are either polynomial or regular images of some Euclidean space. The positive answer to the open quadrant problem has been a recurrent starting point for this approach.

In [4, 5] we presented the first steps to approach Problem 1.1. A complete solution to Problem 1.1 for the one-dimensional case appears in [3], whereas in [6, 8, 9, 17, 18] we approached constructive results concerning the representation as either polynomial or regular images of the semialgebraic sets with piecewise linear boundary commented

above. Articles [7, 10] are of different nature because we find in them new obstructions for a subset of \mathbb{R}^m to be either a polynomial or a regular image of \mathbb{R}^n . In the first one we found some properties of the difference $\text{Cl}(S) \setminus S$ while in the second it is shown that *the set of points at infinite of a polynomial image of \mathbb{R}^n is a connected set.*

The constructive solution to the open quadrant problem provided in [4] involves quite complicated computer calculations that the third author never liked. In fact he provided in his Ph.D. Thesis a different topological proof for the map proposed in [4], together with an algebraic proof involving a different polynomial map. This map has inspired the first and third authors for a short algebraic proof of the open quadrant problem involving a new polynomial map [11] and has led us to look for a polynomial map with optimal algebraic structure whose image is the open quadrant. It is important to establish clearly the meaning of ‘optimal algebraic structure’ [11, §3(A)]. It is natural to wonder how a polynomial map looks like when completely expanded and how it compares with other polynomial maps. We care about the total degree of the involved polynomial map (the sum of the degrees of its components) and its total number of (non-zero) monomials. We would like to find a polynomial map with the less possible total degree and the less possible number of monomials. The example in [4] has total degree 56 and its total number of monomials is 168. The polynomial map in [11] has total degree 72 and its total number of monomials is 350. In this work we will prove:

Theorem 1.3 *The open quadrant \mathcal{Q} is the image of the polynomial map*

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}^2, (x, y) \mapsto ((x^2y^4 + x^4y^2 - y^2 - 1)^2 + x^6y^4, (x^6y^2 + x^2y^2 - x^2 - 1)^2 + x^6y^4).$$

This polynomial map has total degree 28 and its total number of monomials is 22, which certainly improves the already known explicit solutions to the open quadrant problem. It has been constructed following a similar strategy to that in [4, §3]. Our experience approaching this problem suggests us that this map is surely close to have the optimal desired algebraic structure.

The article is organized as follows. In Section 2 we present all basic notions and topological preliminaries used in Section 3 to prove Theorem 1.3.

2. Topological preliminaries

Denote the closed disc of center the origin and radius $A > 0$ of the plane \mathbb{R}^2 with \mathbb{D}_A . A *warped disc* is a subset $\mathcal{D}_{A,\xi} := \{z = \xi(x, y), x^2 + y^2 \leq A^2\} \subset \mathbb{R}^3$ where $\xi : \mathbb{R}^2 \rightarrow \mathbb{R}$ is a continuous function. Consider the homeomorphism

$$\zeta : \mathbb{R}^3 \rightarrow \mathbb{R}^3, (x, y, z) \mapsto (x, y, z - \xi(x, y))$$

that maps $\mathcal{D}_{A,\xi}$ onto $\mathbb{D}_A \times \{0\}$. The image of $\mathcal{D}_{A,\xi}$ under a permutation of the variables of \mathbb{R}^3 will be also called a warped disc.




Figure 1: The homeomorphism ζ for $\xi(x, y) := \sqrt{B^2 - \min(y^2, B^2)}$ acting on \mathbb{R}^3 .

For each $\varepsilon > 0$ consider the open neighborhood

$$\mathbb{D}_A(\varepsilon) := \{x^2 + y^2 < (A + \varepsilon)^2\} \times (-\varepsilon, \varepsilon) \subset \mathbb{R}^3$$

of \mathbb{D}_A . Clearly, $\mathcal{D}_{A,\xi}(\varepsilon) := \zeta^{-1}(\mathbb{D}_A(\varepsilon))$ is an open neighborhood of $\mathcal{D}_{A,\xi}$ in \mathbb{R}^3 .

Definition 2.1 A (continuous) path $\alpha : [a, b] \rightarrow \mathbb{R}^3$ *meets transversally once the warped disc* $\mathcal{D}_{A,\xi}$ if there exist $s_0 \in (a, b)$ and $\varepsilon > 0$ such that $J := \alpha^{-1}(\mathcal{D}_{A,\xi}(\varepsilon)) = (s_0 - \varepsilon, s_0 + \varepsilon)$ is an open subinterval of $[a, b]$ and $(\zeta \circ \alpha)|_J(t) = (0, 0, t - s_0)$.

Remark 2.2 If the path $\alpha : [a, b] \rightarrow \mathbb{R}^3$ meets transversally once the warped disc $\mathcal{D}_{A,\xi}$, then $\alpha([a, b]) \cap \partial\mathcal{D}_{A,\xi} = \emptyset$.

Let C be a topological space homeomorphic to a closed disc and let $\phi : C \rightarrow \mathbb{R}^3$ be a continuous map. The restriction $\partial\phi := \phi|_{\partial C}$ is called the *boundary map* of ϕ . We say that the boundary map $\partial\phi$ *meets transversally once a warped disc* $\mathcal{D}_{A,\xi} \subset \mathbb{R}^3$ if there exists a parameterization β of ∂C such that $\alpha := \phi \circ \beta$ meets transversally once the warped disc $\mathcal{D}_{A,\xi}$.

Given a path-connected topological space X and a point $x_0 \in X$ we denote the fundamental group of X at the base point x_0 with $\pi_1(X, x_0)$. Each path α starting and ending at x_0 is called a loop with base point x_0 and represents an element of $\pi_1(X, x_0)$, that we denote with $[\alpha]$.

Lemma 2.3 *Let $\mathcal{D}_{A,\xi}$ be a warped disc of \mathbb{R}^3 and let $X := \mathbb{R}^3 \setminus \partial\mathcal{D}_{A,\xi}$. Let $\alpha : [a, b] \rightarrow X$ be a loop with base point $x_0 \in X$ that meets transversally once $\mathcal{D}_{A,\xi}$. Then $[\alpha]$ is a generator of $\pi_1(X, x_0) \cong \mathbb{Z}$.*

Proof. Keep the notations introduced above. Let $s_0 \in (a, b)$ and $\varepsilon > 0$ be such that

$$J := \alpha^{-1}(\mathcal{D}_{A,\xi}(\varepsilon)) = (s_0 - \varepsilon, s_0 + \varepsilon)$$

is an open subinterval of $[a, b]$ and $(\zeta \circ \alpha)|_J(t) = (0, 0, t - s_0)$. After a reparameterization of α we may assume $s_0 = 0$.

As ζ is a homeomorphism of \mathbb{R}^3 , we will prove the statement for $\beta := \zeta \circ \alpha$, $Y := \mathbb{R}^3 \setminus \partial\mathbb{D}_A$ and the base point $y_0 := \beta(-\varepsilon) = (0, 0, -\varepsilon)$. Consider the path $\gamma : [0, 1] \rightarrow \mathbb{R}^3$ given by

$$\gamma(t) := \begin{cases} (3(A + \varepsilon)t, 0, \varepsilon) & \text{if } 0 \leq t \leq \frac{1}{3}, \\ (A + \varepsilon, 0, \varepsilon - (t - \frac{1}{3})6\varepsilon) & \text{if } \frac{1}{3} < t \leq \frac{2}{3}, \\ (A + \varepsilon - 3(A + \varepsilon)(t - \frac{2}{3}), 0, -\varepsilon) & \text{if } \frac{2}{3} < t \leq 1. \end{cases}$$




Figure 2: The path β meets transversally once the disk \mathbb{D}_A .

Write $\beta_0 := \beta|_J$ and $\beta_1 := \beta|_{[\varepsilon, b]} * \beta|_{[a, -\varepsilon]}$. We claim:

$$[\beta] = [\beta_0 * \beta_1] = [\beta_0 * \gamma] \cdot [\gamma^{-1} * \beta_1] = g \cdot e = g,$$

where e and g are respectively the identity element and a generator of $\pi_1(Y, y_0) \cong \mathbb{Z}$.

The loop $\gamma^{-1} * \beta_1$ with base point y_0 is contained in $\mathbb{R}^3 \setminus \mathbb{D}_A$, which is a simply connected space. Consequently, $[\gamma^{-1} * \beta_1] = e$ in $\pi_1(Y, y_0)$.

The class $[\beta_0 * \gamma]$ generates $\pi_1(Y, y_0)$. Indeed, Y has as deformation retract the set $Z := \partial\mathbb{D}_A(\varepsilon) \cup I_\varepsilon$ where $I_\varepsilon := \{(0, 0)\} \times \{-\varepsilon \leq z \leq \varepsilon\}$. It is an exercise of algebraic topology to show that $[\beta_0 * \gamma]$ is a generator of $\pi_1(Z, y_0) \cong \pi_1(Y, y_0) \cong \mathbb{Z}$, as required. \square

Lemma 2.4 *Let $\phi : C \rightarrow X$ be a continuous map and assume that C is homeomorphic to a closed disc. Let $\beta : [a, b] \rightarrow \partial C$ be a parameterization starting and ending at $z_0 \in \partial C$. Then $[\phi \circ \beta]$ is the identity element of $\pi_1(X, \phi(z_0))$.*

Proof. Let $\psi : C \rightarrow \{x^2 + y^2 \leq 1\}$ be a homeomorphism. The continuous map

$$H : [0, 1] \times [a, b] \rightarrow X, (\rho, t) \mapsto (\phi \circ \psi^{-1})(\rho \cdot (\psi \circ \beta)(t) + (1 - \rho) \cdot \psi(z_0))$$

is a homotopy map between $\phi \circ \beta$ and the constant path, as required. \square

Proposition 2.5 *Let C be a topological space homeomorphic to a closed disc and $\phi : C \rightarrow \mathbb{R}^3$ a continuous map. Assume $\partial\phi : \partial C \rightarrow \mathbb{R}^3$ meets transversally once a warped disc $\mathcal{D} \subset \mathbb{R}^3$. Then $\partial\mathcal{D} \cap \phi(\text{Int}(C)) \neq \emptyset$.*

Proof. Assume by contradiction $\partial\mathcal{D} \cap \phi(\text{Int}(C)) = \emptyset$. As $\partial\phi$ meets transversally once \mathcal{D} , the image $\phi(\partial C)$ does not intersect $\partial\mathcal{D}$ by Remark 2.2. Thus, $\phi(C) \subset X := \mathbb{R}^3 \setminus \partial\mathcal{D}$. Let $\beta : [a, b] \rightarrow \partial C$ be a parameterization starting and ending at $z_0 \in \partial C$ such that $\phi \circ \beta$ meets transversally once \mathcal{D} . By Lemma 2.4 the class $[\phi \circ \beta]$ is the identity element of $\pi_1(X, \phi(z_0))$. However, by Lemma 2.3 the class $[\phi \circ \beta]$ is a generator of $\pi_1(X, \phi(z_0)) \cong \mathbb{Z}$, which is a contradiction. Consequently, $\partial\mathcal{D} \cap \phi(\text{Int}(C)) \neq \emptyset$, as required. \square

3. Proof of Theorem 1.3

Observe first that the map f in the statement of Theorem 1.3 is the composition $f_2 \circ f_1$ of the polynomial maps

$$\begin{aligned} f_1 : \mathbb{R}^2 &\rightarrow \mathbb{R}^2, & (x, y) &\mapsto (x^2, y^2), \\ f_2 : \mathbb{R}^2 &\rightarrow \mathbb{R}^2, & (x, y) &\mapsto ((xy^2 + x^2y - y - 1)^2 + x^3y^2, (x^3y + xy - x - 1)^2 + x^3y^2). \end{aligned}$$

As $f_1(\mathbb{R}^2)$ is the closed quadrant $\bar{\mathcal{Q}} := \{x \geq 0, y \geq 0\}$, we have to prove the equality

$$f_2(\bar{\mathcal{Q}}) = \mathcal{Q}. \tag{3.1}$$

The inclusion $f_2(\overline{\mathcal{Q}}) \subset \mathcal{Q}$ is straightforward because both components of f_2 are strictly positive on $\overline{\mathcal{Q}}$. It only remains to show the inclusion

$$\mathcal{Q} \subset f_2(\overline{\mathcal{Q}}). \tag{3.2}$$

3.1. Reduction of the proof of inclusion (3.2)

Consider the (continuous) semialgebraic maps

$$\begin{aligned} g : \overline{\mathcal{Q}} &\rightarrow \mathbb{R}^3, (x, y) \mapsto (xy^2 + x^2y - y - 1, x^{3/2}y, x^3y + xy - x - 1) \\ h : \mathbb{R}^3 &\rightarrow \mathbb{R}^2, (x, y, z) \mapsto (x^2 + y^2, y^2 + z^2). \end{aligned}$$

As $f_2 = h \circ g$, we have to show that for each tuple $(A^2, B^2) \in \mathcal{Q}$ there exists $(x_0, y_0) \in \overline{\mathcal{Q}}$ such that $(h \circ g)(x_0, y_0) = (A^2, B^2)$. This is equivalent to check that the intersection $h^{-1}(\{(A^2, B^2)\}) \cap g(\overline{\mathcal{Q}})$ is non-empty.

Denote $\mathcal{S} := g(\overline{\mathcal{Q}})$ and fix values $B \geq A > 0$. It holds that sets

$$\begin{aligned} h^{-1}(\{(A^2, B^2)\}) &= \{x^2 + y^2 = A^2, y^2 + z^2 = B^2\}, \\ h^{-1}(\{(B^2, A^2)\}) &= \{y^2 + z^2 = A^2, x^2 + y^2 = B^2\} \end{aligned}$$

contain respectively the boundaries of the warped discs

$$\mathcal{D}_1 : z = \xi_1(x, y), \quad x^2 + y^2 \leq A^2, \tag{3.3}$$

$$\mathcal{D}_2 : x = \xi_2(y, z), \quad y^2 + z^2 \leq A^2, \tag{3.4}$$

for the (continuous) semialgebraic functions

$$\xi_1 : \mathbb{R}^2 \rightarrow \mathbb{R}, (x, y) \mapsto \sqrt{B^2 - \min\{y^2, B^2\}}, \tag{3.5}$$

$$\xi_2 : \mathbb{R}^2 \rightarrow \mathbb{R}, (y, z) \mapsto \sqrt{B^2 - \min\{y^2, B^2\}}. \tag{3.6}$$

Consequently, we are reduced to prove:

3.1.1. *For fixed values $B \geq A > 0$ the intersections $\partial\mathcal{D}_1 \cap \mathcal{S}$ and $\partial\mathcal{D}_2 \cap \mathcal{S}$ are non-empty.*

3.2. Proof of Statement 3.1.1

Write $\mathcal{R} := [0, +\infty) \times (0, \frac{\pi}{2})$ and $\overline{\mathcal{R}} := [0, +\infty) \times [0, \frac{\pi}{2}]$. Consider the map $\phi := (\phi_1, \phi_2, \phi_3) : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ where

$$\begin{aligned} \phi_1(\rho, \theta) &:= \cos \theta \sin \theta (\cos \theta - \sin \theta)^2 \\ &\quad + \rho(2 \cos^4 \theta \sin \theta + \cos \theta \sin^4 \theta + \cos^5 \theta) + \rho^2 \cos^5 \theta \sin \theta, \\ \phi_2(\rho, \theta) &:= \sqrt{\cos \theta \sin \theta} (\cos \theta + \sin \theta + \rho \cos \theta \sin \theta), \\ \phi_3(\rho, \theta) &:= \rho \sin \theta. \end{aligned}$$

Let us prove now some properties of the map ϕ and the sets \mathcal{R} and $\overline{\mathcal{R}}$:

3.2.1. $\phi(\mathcal{R}) \subset \mathcal{S}$.

Proof. The analytic map

$$\psi : \mathcal{R} \rightarrow \mathcal{Q}, \quad (\rho, \theta) \mapsto \left(\frac{\sin \theta}{\cos \theta}, \frac{(\cos \theta + \sin \theta + \rho \cos \theta \sin \theta) \cos^2 \theta}{\sin \theta} \right),$$

satisfies $\psi(\mathcal{R}) \subset \overline{\mathcal{Q}}$ and $g \circ \psi = \phi|_{\mathcal{R}}$. Consequently, $\phi(\mathcal{R}) \subset \mathcal{S}$, as required. \square

3.2.2. The inequality $\phi_1^2(\rho, \theta) + \phi_3^2(\rho, \theta) \geq \frac{\rho^2}{4}$ holds for each $(\rho, \theta) \in \overline{\mathcal{R}}$. Consequently,

$$\text{dist}(\phi(\rho, \theta), \mathbf{0}) \geq \frac{\rho}{2} \tag{3.7}$$

for each $(\rho, \theta) \in \overline{\mathcal{R}}$.

Proof. As $\rho, \cos \theta, \sin \theta$ are ≥ 0 on $\overline{\mathcal{R}}$, we have

$$\begin{aligned} \phi_1(\rho, \theta) &\geq \rho \cos \theta (\cos^4 \theta + \sin^4 \theta) = \rho \cos \theta (1 - 2 \cos^2 \theta \sin^2 \theta) \\ &= \rho \cos \theta \left(1 - \frac{\sin^2(2\theta)}{2} \right) \geq \frac{\rho}{2} \cos \theta. \end{aligned}$$

In addition, $\phi_3(\rho, \theta) = \rho \sin \theta \geq \frac{\rho}{2} \sin \theta$, so

$$\phi_1^2(\rho, \theta) + \phi_3^2(\rho, \theta) \geq \frac{\rho^2}{4} \cos^2 \theta + \frac{\rho^2}{4} \sin^2 \theta = \frac{\rho^2}{4},$$

as required. \square

3.2.3. The map ϕ satisfies $\phi(0, \theta) = \phi(0, \frac{\pi}{2} - \theta)$ for $\theta \in [0, \frac{\pi}{2}]$. Fix $M > 0$ and consider the rectangle $\overline{\mathcal{R}}_M := [0, M] \times [0, \frac{\pi}{2}]$. Denote $\phi_M := \phi|_{\overline{\mathcal{R}}_M}$. Identify the points $(0, \theta)$ and $(0, \frac{\pi}{2} - \theta)$ for $\theta \in [0, \frac{\pi}{2}]$ and endow the quotient space $\overline{\tilde{\mathcal{R}}}_M$ with the quotient topology. Observe that the interior $\text{Int}(\overline{\tilde{\mathcal{R}}}_M)$ of $\overline{\tilde{\mathcal{R}}}_M$ as a topological manifold with boundary is the quotient space $\tilde{\mathcal{R}}_M$ obtained identifying the points $(0, \theta)$ and $(0, \frac{\pi}{2} - \theta)$ of $\mathcal{R}_M := [0, M] \times (0, \frac{\pi}{2})$, where $\theta \in (0, \frac{\pi}{2})$.

The canonical projection $\pi_M : \overline{\mathcal{R}}_M \rightarrow \overline{\tilde{\mathcal{R}}}_M$ is continuous. As ϕ_M is compatible with π_M , there exists a continuous map $\tilde{\phi}_M : \overline{\tilde{\mathcal{R}}}_M \rightarrow \mathbb{R}^3$ such that the following diagram is commutative. In addition, $\tilde{\phi}_M(\tilde{\mathcal{R}}_M) = \phi(\mathcal{R}_M) \subset \mathcal{S}$.

$$\begin{array}{ccc} \mathcal{R}_M & \hookrightarrow & \overline{\mathcal{R}}_M \\ \pi_M|_{\mathcal{R}_M} \downarrow & & \downarrow \pi_M \quad \searrow \phi_M \\ \tilde{\mathcal{R}}_M & \hookrightarrow & \overline{\tilde{\mathcal{R}}}_M \xrightarrow{\tilde{\phi}_M} \mathbb{R}^3 \end{array}$$




Figure 3: Left and right views of $\phi_M(\mathcal{R}_M) \subset \mathcal{S}$.

3.2.4. $\tilde{\mathcal{R}}_M$ is homeomorphic to a disc and its boundary is the set

$$\pi_M(\{\rho = M\} \cup \{\theta = 0\} \cup \{\theta = \frac{\pi}{2}\}).$$

Proof. Identify \mathbb{R}^2 with \mathbb{C} (interchanging the order of the variables $(\rho, \theta) \rightsquigarrow (\theta, \rho)$) and consider the continuous map

$$\mu : \mathbb{C} \rightarrow \mathbb{C}, \quad z := \theta + \sqrt{-1}\rho \mapsto w := u + \sqrt{-1}v = \left(\frac{4}{\pi}z - 1\right)^2.$$

The restriction $\mu|_{\{\rho > 0\}} : \{\rho > 0\} \rightarrow \mathbb{C} \setminus ([0, +\infty) \times \{0\})$ is a homeomorphism and the image of $\bar{\mathcal{R}}_M \setminus \{\rho = 0\}$ is

$$\mathcal{T}_M := \{(u, v) \in \mathbb{R}^2 : (\frac{\pi v}{8M})^2 - (\frac{4M}{\pi})^2 \leq u \leq 1 - (\frac{v}{2})^2\} \setminus ([0, 1] \times \{0\}).$$

The closure $\bar{\mathcal{T}}_M$ of \mathcal{T}_M is a compact convex set (as it is a closed bounded intersection of two convex sets). By [1, Cor.11.3.4] $\bar{\mathcal{T}}_M$ is homeomorphic to a closed disc. In addition

$$\mu|_{\{\rho=0\}} : \{\rho = 0\} \rightarrow [0, +\infty) \times \{0\}, \quad \theta \mapsto \left(\frac{4}{\pi}\theta - 1\right)^2$$

transforms the segment $[0, \frac{\pi}{2}] \times \{0\}$ onto the interval $[0, 1]$. The preimage of $t_0 \in [0, 1]$ under $\mu|_{\{\rho=0\}}$ is

$$\{\theta_1 := \frac{\pi}{4}(1 + \sqrt{t_0}), \theta_2 := \frac{\pi}{4}(1 - \sqrt{t_0})\}.$$

As $\theta_1 = \frac{\pi}{2} - \theta_2$, the map $\lambda := \mu|_{\bar{\mathcal{R}}_M} : \bar{\mathcal{R}}_M \rightarrow \bar{\mathcal{T}}_M$ factors through $\tilde{\mathcal{R}}_M$ and there exists a continuous map $\tilde{\lambda} : \tilde{\mathcal{R}}_M \rightarrow \bar{\mathcal{T}}_M$ such that the following diagram is commutative.




Figure 4: Behavior of the map $\mu : \bar{\mathcal{R}}_M \rightarrow \bar{\mathcal{T}}_M$.

$$\begin{array}{ccccc}
 \mathcal{R}_M & \hookrightarrow & \bar{\mathcal{R}}_M & & \\
 \pi_M|_{\mathcal{R}_M} \downarrow & & \pi_M \downarrow & \searrow \lambda & \\
 \tilde{\mathcal{R}}_M & \hookrightarrow & \tilde{\bar{\mathcal{R}}}_M & \xrightarrow{\tilde{\lambda}} & \mathcal{S}_M
 \end{array}$$

The map $\tilde{\lambda}$ is continuous and bijective and it maps the compact set $\tilde{\bar{\mathcal{R}}}_M$ onto the Hausdorff space $\bar{\mathcal{T}}_M$, so it is a homeomorphism. Consequently, $\tilde{\bar{\mathcal{R}}}_M$ is homeomorphic to a disc and its boundary is $\pi_M(\{\rho = M\} \cup \{\theta = 0\} \cup \{\theta = \frac{\pi}{2}\})$, as required. \square

3.2.5. Fix $B \geq A > 0$ and consider the warped discs \mathcal{D}_1 and \mathcal{D}_2 introduced in (3.3) and (3.4). Then there exists $M > 0$ such that the boundary map $\partial\tilde{\phi}_M : \partial\tilde{\mathcal{R}}_M \rightarrow \mathbb{R}^3$ meets transversally once both discs \mathcal{D}_1 and \mathcal{D}_2 .

Proof. As \mathcal{D}_1 and \mathcal{D}_2 are bounded set, there exists $M_0 > 0$ such that $\mathcal{D}_1 \cup \mathcal{D}_2 \subset \{\|(x, y, z)\| < M_0\}$. Take $M := 4M_0$ and consider the set $\tilde{\bar{\mathcal{R}}}_M$ and the continuous map $\tilde{\phi}_M$ introduced in paragraph 3.2.3.

We claim: the boundary map $\partial\tilde{\phi}_M : \partial\tilde{\mathcal{R}}_M \rightarrow \mathbb{R}^3$ meets transversally once \mathcal{D}_1 .




Figure 5: The boundary map $\partial\tilde{\phi}_M : \partial\tilde{\mathcal{R}}_M \rightarrow \mathbb{R}^3$ meets transversally once \mathcal{D}_1 .

Consider the parameterization of $\partial\tilde{\mathcal{R}}_M$ given by

$$\beta_1(t) := \begin{cases} \pi_M(t, \frac{\pi}{2}), & \text{if } 0 \leq t \leq M, \\ \pi_M(M, M + \frac{\pi}{2} - t), & \text{if } M < t \leq M + \frac{\pi}{2}, \\ \pi_M(2M + \frac{\pi}{2} - t, 0), & \text{if } M + \frac{\pi}{2} < t \leq 2M + \frac{\pi}{2}. \end{cases}$$

We have

$$\alpha_1(t) := \tilde{\phi}_M \circ \beta_1(t) = \begin{cases} \phi(t, \frac{\pi}{2}), & \text{if } 0 \leq t \leq M, \\ \phi(M, M + \frac{\pi}{2} - t), & \text{if } M < t \leq M + \frac{\pi}{2}, \\ \phi(2M + \frac{\pi}{2} - t, 0), & \text{if } M + \frac{\pi}{2} < t \leq 2M + \frac{\pi}{2}. \end{cases}$$

Choose $0 < \varepsilon < \min\{B, M_0 - B\}$ and consider the homeomorphism

$$\zeta_1 : \mathbb{R}^3 \rightarrow \mathbb{R}^3, (x, y, z) \mapsto (x, y, z - \xi_1(x, y)),$$

where ξ_1 is the (continuous) semialgebraic function introduced in (3.5). Denote $\mathcal{D}_1(\varepsilon) := \zeta_1^{-1}(\mathbb{D}_A(\varepsilon))$. It is enough to check:

$$\alpha_1^{-1}(\mathcal{D}_1(\varepsilon)) = (B - \varepsilon, B + \varepsilon).$$

Pick $p_0 := \alpha_1(t_0) \in \text{Im}(\alpha_1)$. We distinguish three cases:

- (i) If $0 \leq t_0 \leq M$, then $\zeta_1(p_0) = (\zeta_1 \circ \phi)(t_0, 0) = (0, 0, t_0 - B)$. Consequently, $\zeta_1(p_0) \in \mathbb{D}_A(\varepsilon)$ if and only if $-B < -\varepsilon < t_0 - B < \varepsilon < M - B$.




Figure 6: Behavior of the paths β_1 and β_2 .

(ii) If $M < t_0 \leq M + \frac{\pi}{2}$, we have by (3.7)

$$\text{dist}(p_0, \mathbf{0}) \geq \frac{M}{2} = 2M_0 > \sqrt{2}M_0 > \text{dist}(q, \mathbf{0})$$

for each $q \in \mathcal{D}_1(\varepsilon)$. Therefore $p_0 \notin \mathcal{D}_1(\varepsilon)$.

(iii) If $M + \frac{\pi}{2} < t_0 \leq 2M + \frac{\pi}{2}$, then

$$p_0 = \alpha_1(t_0) = \phi(2M + \frac{\pi}{2} - t_0, 0) = (2M + \frac{\pi}{2} - t_0, 0, 0),$$

so $\zeta_1(p_0) = (2M + \frac{\pi}{2} - t_0, 0, -B)$. As $\varepsilon < B$, it holds $\zeta_1(p_0) \notin \mathbb{D}_A(\varepsilon)$, so $p_0 \notin \mathcal{D}_1(\varepsilon)$.

We conclude $\alpha_1^{-1}(\mathcal{D}_1(\varepsilon)) = (B - \varepsilon, B + \varepsilon)$, so α_1 meets transversally once \mathcal{D}_1 .

Analogously one shows: *the boundary map $\partial\tilde{\mathcal{R}}_M : \partial\tilde{\mathcal{R}}_M \rightarrow \mathbb{R}^3$ meets transversally once \mathcal{D}_2 .*

Consider in this case the parameterization of $\partial\tilde{\mathcal{R}}_M$ given by

$$\beta_2(t) := \begin{cases} \pi_M(t, 0), & \text{if } 0 \leq t \leq M, \\ \pi_M(M, t - M), & \text{if } M < t \leq M + \frac{\pi}{2}, \\ \pi_M(2M + \frac{\pi}{2} - t, \frac{\pi}{2}), & \text{if } M + \frac{\pi}{2} < t \leq 2M + \frac{\pi}{2}. \end{cases}$$

We have

$$\alpha_2(t) := \tilde{\phi}_M \circ \beta_2(t) = \begin{cases} \phi(t, 0), & \text{if } 0 \leq t \leq M, \\ \phi(M, t - M), & \text{if } M < t \leq M + \frac{\pi}{2}, \\ \phi(2M + \frac{\pi}{2} - t, \frac{\pi}{2}), & \text{if } M + \frac{\pi}{2} < t \leq 2M + \frac{\pi}{2}. \end{cases}$$




Figure 7: The boundary map $\partial\tilde{\phi}_M : \partial\tilde{\mathcal{R}}_M \rightarrow \mathbb{R}^3$ meets transversally once \mathcal{D}_2 .

Proceed as above keeping the same values for A and ε and using in this case the homeomorphism

$$\zeta_2 : \mathbb{R}^3 \rightarrow \mathbb{R}^3, (x, y, z) \mapsto (z, y, x - \xi_2(z, y)),$$

where ξ_2 is the (continuous) semialgebraic function introduced in (3.6), to prove that α_2 meets transversally once the warped disk \mathcal{D}_2 . \square

3.2.6. By 3.2.4 $\tilde{\mathcal{R}}_M$ is homeomorphic to a closed disc. By Proposition 2.5 applied to the continuous map $\tilde{\phi}_M : \tilde{\mathcal{R}}_M \rightarrow \mathbb{R}^3$ and 3.2.5, we deduce that the boundaries of both warped discs \mathcal{D}_1 and \mathcal{D}_2 meet $\phi_M(\mathcal{R}_M) \subset \mathcal{S}$. Thus, 3.1.1 holds, as required. \square

References

- [1] M. Berger: *Geometry, I*. Universitext, Springer-Verlag, Berlin 1987.
- [2] J. F. Fernando: *Polynomial images of \mathbb{R}^n* . In *Reelle algebraische und analytische Geometrie, March 17–23 (2002)* Oberwolfach.
- [3] J. F. Fernando: *On the one-dimensional polynomial and regular images of \mathbb{R}^n* . *J. Pure Appl. Algebra* **218**, 9(2014) 1745–1753.
- [4] J. F. Fernando, J. M. Gamboa: *Polynomial images of \mathbb{R}^n* . *J. Pure Appl. Algebra* **179**, 3 (2003) 241–254.
- [5] J. F. Fernando, J. M. Gamboa: *Polynomial and regular images of \mathbb{R}^n* . *Israel J. Math.* **153** (2006) 61–92.

- [6] J. F. Fernando, J. M. Gamboa, C. Ueno: *On convex polyhedra as regular images of \mathbb{R}^n* . Proc. London Math. Soc. (3) **103** (2011) 847–878.
- [7] J. F. Fernando, J. M. Gamboa, C. Ueno: *Sobre las propiedades de la frontera exterior de las imágenes polinómicas y regulares de \mathbb{R}^n* . In *Contribuciones Matemáticas en homenaje a Juan Tarrés*, 159–178, UCM 2012.
- [8] J. F. Fernando, C. Ueno: *On complements of convex polyhedra as polynomial and regular images of \mathbb{R}^n* . Int. Math. Res. Not. **18** (2014) 5084–5123.
- [9] J. F. Fernando, C. Ueno: *On the complements of 3-dimensional convex polyhedra as polynomial images of \mathbb{R}^3* . Internat. J. Math. **25**, 7 (2014) 1450071, 18pp.
- [10] J. F. Fernando, C. Ueno: *On the set of points at infinity of a polynomial image of \mathbb{R}^n* . Discrete Comput. Geom. **52**, 4 (2014) 583–611.
- [11] J. F. Fernando, C. Ueno: *A short proof for the open quadrant problem*. Preprint RAAG (2014, submitted to MEGA 2015) 8 pp.
- [12] J. M. Gamboa: *Algebraic images of the real plane*. In *Reelle algebraische Geometrie, June 10-16 (1990)* Oberwolfach.
- [13] A. Harnack: *Über die Vielheiligkeit der ebenen algebraischen Kurven*. Math. Ann. **10** (1876) 189–198.
- [14] S. Pinchuk: *A counterexample to the real Jacobian Conjecture*. Math. Z. **217** (1994) 1–4.
- [15] A. Tarski: *A decision method for elementary algebra and geometry*. Prepared for publication by J.C.C. Mac Kinsey, Berkeley 1951.
- [16] C. Ueno: *Imágenes polinómicas y regulares de espacios euclídeos*. Ph.D. Thesis UCM 2012.
- [17] C. Ueno: *A note on boundaries of open polynomial images of \mathbb{R}^2* . Rev. Mat. Iberoam. **24**, 3 (2008) 981–988.
- [18] C. Ueno: *On convex polygons and their complements as images of regular and polynomial maps of \mathbb{R}^2* . J. Pure Appl. Algebra **216**, 11 (2012) 2436–2448.